



International Journal of Virology & Infectious Diseases

Review Article

A Review on Knowledge, Attitude, and Practice during the COVID-19 Pandemic in Ethiopia -

Samuel Chane Teferi*

Department of Biology, Salale University, Ethiopia

***Address for Correspondence:** Samuel Chane Teferi, Department of Biology, Salale University, Ethiopia, ORCID: <https://orcid.org/0000-0002-5805-7857>; E-mail: Samuelchane19@gmail.com

Submitted: 20 August 2020; Approved: 25 August 2020; Published: 29 August 2020

Cite this article: Teferi SC. A Review on Knowledge, Attitude, and Practice during the COVID-19 Pandemic in Ethiopia. *Int J Virol Infect Dis.* 2020;5(2): 040-044.

Copyright: © 2020 Teferi SC. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



ABSTRACT

World Health Organization declared the novel coronavirus as a global public health emergency on January 30, 2020. COVID-19 has infected more than 17,660,523 people worldwide, with more than 680,894 deaths in different regions and countries. Based on the World Health Organization (WHO) Africa report, as of 02 August, 2020, a cumulative total of 802,792 confirmed COVID-19 cases with 13,779 deaths have been reported across all African countries in the region. In Ethiopia the virus spreads alarmingly because the community didn't practice the information given by Ministry of Health and the Government; as of August 2, 2020, 17,999 cases and 284 deaths. The level of panic in peoples is associated with knowledge and attitude according to Researchers. Good Knowledge, Attitude and Practice can be used to assess the current conditions and if it is used properly it plays a pivotal role to control the spread of COVID-19. Therefore, the government and ministry of health must give information to the society through TV, radio and social media repeatedly. So, the aim of the current study was to assess/review the knowledge, attitude and perception of the population towards COVID-19 pandemic in Ethiopia.

Keywords: Ethiopia; Practice; COVID-19; Pandemic

INTRODUCTION

The 2019 novel coronavirus (2019-nCoV) or the severe acute respiratory syndrome corona virus 2 (SARS-CoV-2) as it is now called, is rapidly spreading from its origin in Wuhan City of Hubei Province of China to the rest of the world [1]. Novel coronavirus (pandemic) was named as "Corona Virus Disease 2019" (2019-nCoV) by World Health Organization (WHO) in Geneva, Switzerland [2,3]. So this virus is deadly third-generation virus in Corona family apart from Middle East Respiratory Syndrome (MERS) in 2012 and Severe Acute Respiratory Syndrome (SARS) in 2003. The diameter of corona virus is 65-125 nm, and contain a single strand of RNA with lengths ranging from 26 to 32 kb and it is under Coronaviridae family. Coronaviruses comprise several types, such as alpha, beta, gamma, delta, SARSCoV, H5N1 influenza A, H1N1 2009, and MERS-CoV [4].

World Health Organization declared the novel corona virus as a global public health emergency (pandemic) on January 30, 2020 [5]. Novel coronavirus disease is a highly infectious disease, and the ongoing outbreak has affected a huge part of populations around the world. There are four levels of COVID-19 transmission according to World Health Organization, such as, no cases reported, sporadic cases, Clusters of cases, or Community transmission [6]. Public health and social measures play a crucial role in reducing the number of infections and reduce death until the vaccine is ready [1].

The novel corona virus is transmitted through large droplets generated during coughing and sneezing by symptomatic and sometimes from asymptomatic patients [7]. Therefore, frequent hand-washing with soap and water and using sanitizer or alcohol is crucial. The common clinical features include fever (not in all), cough, sore throat, headache, fatigue, headache, myalgia and breathlessness [8].

COVID-19 has infected more than 17,660,523 people worldwide, with more than 680,894 deaths in different regions and countries. The USA, the major hit country alone, recorded more than 359,180 deaths on early August 2020. Based on the World Health Organization (WHO) Africa report, as of 02 August, 2020, a cumulative total of 802,792 confirmed COVID-19 cases with 13,779 deaths have been reported across all African countries in the region [9]. In Ethiopia the virus spreads alarmingly because the community didn't practice the information given by ministry of health and the government; as of August 2, 2020, 17,999 cases and 284 deaths have been reported [9].

Subsequently, Ethiopia has also commenced monitoring the disease and has implemented the COVID-19 prevention and containment interventions recommended by World Health Organization [10]. Health system of Ethiopia is not as developed as

other countries so if the virus is not contained it will cost the country many things. Knowledge, Attitude, and Practice (KAP) research is important to collect information on what is known, believed, and done by a specific population [11]. But in Ethiopia those types of research is not available like other countries. It is vital to know the level of knowledge, attitude, and practice of the population towards COVID-19 to contain the virus in countries like Ethiopia. Therefore, this review aimed to assess the knowledge, attitude and perception of COVID-19 pandemic in Ethiopia.

LITERATURE SEARCH

A literature search was conducted in August 1-2, 2020 using the keywords "Practice," "Attitude," "Knowledge," "Ethiopia, and "COVID-19" on PubMed and Google Scholar databases and the reference list of all identified reports and articles were searched manually in Google. The search yields a total of 13 articles.

KAP STUDIES ON COVID-19

Thirteen articles on KAP during COVID-19 in Ethiopia (Table 1). The articles used questionnaires and interview (including face to face and telephone) for data collection and the number of respondents ranging from 247 to 1570 for a total of 7,335. The respondents consisted of health workers, population, health students and patients from different corners of the country.

KNOWLEDGE

Thirteen articles on knowledge about COVID-19 among health workers, hospital staff, students, and sample populations. All articles reported substantial levels of knowledge about COVID-19 in Ethiopia. A research on 422 health workers showed that nurses obtained higher knowledge scores which is very important to tackle the virus because nurses are one of the frontline workers.

In a study by [12] the majority (70.1%) of the study participants reported that shaking hands of infected individuals result in the spread of infection. 217 (53.7%) knew touching an object or surface with the virus on it, then touching the mouth, nose, or eye transmits the virus and 337 (83.4%) knew respiratory droplets of infected individuals through the air during sneezing or coughing spreads the virus.

In other study, from 546 participants, 71.4% correctly responded that the main clinical symptoms of COVID-19 are fever, fatigue, dry cough, and shortness of breath, and the majority (95%) said currently there is no cure for COVID-19 and more than 73.6% of the participants used social media as their main source of information about COVID-19 [14].



Table 1: List of articles conducted on KAP in Ethiopia.

NO.	Type of study	Participants	Number of participant	Instrument	Author
1	Cross-sectional study on KAP	Patients (chronic diseases)	404	Questionnaire	[12]
2	Cross-sectional survey on KAP	Arba Minch Town (Population)	528	Online questioner	[13]
3	Cross-sectional survey on KAP	Students at Debre Berhan University	546	Questionnaire	[14]
4	Cross-sectional study on KAP	Population	1570	Telephone interview	[15]
5	Cross-sectional study on KAP	Healthcare workers	422	Questionnaire	[16]
6	Cross-sectional survey on KA	Population	1037	phone-based survey	[17]
7	Cross-Sectional Study on KAP	Students (Amhara)	408	Questionnaire	[18]
8	Cross-sectional study on KAP	Nurses	415	Questionnaire	[19]
9	Cross-sectional study on KAP	Patients visiting JMC	247	Interview	[20]
10	Online Cross-Sectional Study KP	Population	341	Online questioner	[21]
11	Online Cross-Sectional Survey on KP	Educated individuals	528	Online questioner	[22]
12	Cross sectional study on AP	Southern Ethiopia population	585	Interview	[23]
13	Cross sectional study on K	Health science students	304	Questioner	[24]
		Total	7,335		

Abbreviations: K = Knowledge; A = Attitude; P = Practice.

Table 2: Knowledge studies.

No	Author	Participant	Instrument	Knowledge score	Knowledge result
1	[12]	Patients (chronic diseases)	Questioner	33.9% (95% CI (29.3–38.5%))	One-third of chronic disease patients had poor knowledge
2	[13]	Arba Minch Town (Population)	Online questioner	11.48 (SD: 2.25, range: 1-15)	The majority of respondents had good knowledge
3	[14]	Students at Debre Berhan University	Questioner	9.6 ± 1.8 with a range of 0–13	Most of participants (403 (73.8%)) considered as having good knowledge.
4	[15]	Population	Telephone interview	4.2 (SD=2.809, range 0-10)	knowledge is unsatisfactory
5	[16]	Healthcare workers	Questionnaire	350 (88.2%)	Good knowledge of healthcare workers
6	[17]	Population	phone-based survey	6.9 (SD:1.65)	There is a good level of knowledge in the population
7	[18]	Students (Amhara)	Questionnaire	284 (69.6%) (95% CI 65% 74.3%)	Overall good knowledge was below the WHO recommendation scores.
8	[19]	Nurses	Questionnaire	307(74% (with CI; 70 to 78.1)	Nurses had good knowledge which is vital to defeat the virus
9	[20]	Patients visiting JMC	Interview	High knowledge 41.3%, Moderate (41.7%), Low (17%)	The visitors' knowledge was modest to protect themselves from this highly contagious virus.
10	[21]	Population	Online questioner	5.52 ± 1.11 Range (0-7)	78.8% had good Knowledge
11	[22]	Educated individuals	Online questioner	(295/528, 55.9%)	Good knowledge
12	[24]	Health science students	Questioner	Good knowledge 25%, poor knowledge 75%	Very poor knowledge in health science students

Respondents heard about COVID-19 were (73.6%), 42.4% and 37.8% knew COVID-19 transmission and clinical manifestations, respectively. Respondents Amhara (AOR = 0.497 (95%CI: 0.2990-0.6542) and Oromia (AOR = 0.4512; 95%CI: 216 0.3218-0.8026) regional states had poor level of knowledge compared with respondents from Addis Ababa according to [15]. A study by [16] revealed that 350 (88.2%) of participants had good knowledge about COVID-19. 75.5% and 82.6% of participants said, that COVID-19 has no specific treatment and vaccine respectively and the majority of healthcare workers gathered information regarding COVID-19 from social media (73.6%) and television (71.5%). However, sources of

knowledge about COVID-19 are TV/Radio has a role of 424(80.3%) and followed by a social-media 372(70.5%) according to [13].

A study by [18] showed that 276 (67.6%) of the students said that air droplets from the infected persons can transmit the infection of COVID-19 to healthy individuals. Similarly, 375 (91.9%), 343 (84.1%), and 324 (79.4%) of the participants said that patients with COVID-19 can present with fever, dry cough, and shortness of breath respectively and 293 (71.8%) of the students have gotten information about COVID-19 from mass media (TV, magazines, newspaper, radio) and nearly fifty percent (54.2%) of the participants



have gotten information from social media (facebook, Instagram, whatsapp and telegram).

ATTITUDE TOWARD COVID-19

According to [13] Knowledge is a prerequisite for establishing prevention beliefs, forming positive attitudes, and promoting positive behaviours, and individuals' cognition and attitudes towards disease affect the effectiveness of their coping strategies and behaviours to a certain extent. A study by [19] showed (72%) of the study participants had favorable attitude towards the COVID-19 and 85.3% of the nurses had disturbed psychological responses towards the COVID-19. The vast majority of the participants also held an optimistic attitude towards the COVID-19 epidemic according to [13]: 81.8% believed that COVID-19 will finally be successfully controlled, and 77.3% had

confidence that world leader/WHO can win the battle against the virus.

PRACTICE TOWARD COVID-19

Table 4 presents articles on the practice of COVID- 19 pandemic in Ethiopia. Two hundred sixty-five (65.5%) study participants reported that they washed their hands with soap frequently, (71.7%) of the respondents had avoided handshaking, (36.6%) of the study participants used face mask, 154 (38.1%) attending overcrowded place, 224 (55.2%) didn't clean frequently touched surfaces and objects and 121 (29.9%) didn't Practice physical distance [12]. Two hundred sixteen (40.9%) of participants gone crowded place and 336(63.3%) were didn't used face-mask when leaving their home. Three hundred ninety-six (75.0%) respondents were used sterilizers before and after touching inanimate object [13].

Table 3: Attitude studies in Ethiopia.

No	Author	Participant	Instrument	Attitude score	Result
1	[12]	Patients (chronic diseases)	Questioner	NA	222 (54.9%), 198 (49.0%), and 71 (17.6%) considered it easy: Avoiding; touching face with the unwashed hand, shaking others, and attending in a crowded population. nearly half of the study participant afraid of contracting the virus.
2	[13]	Arba Minch Town (Population)	Online questioner	NA	82.6% of study participants washed their hands frequently with soap and water
3	[14]	Students at Debre Berhan University	Questioner	NA	229 (42%) they have no concern of being infected with COVID-19.
4	[15]	Population	Telephone interview	NA	50% of the respondents either agree or strongly agree that traditional herbs and religious faith such as holy water can cure COVID-19. Half of the respondents think that it is unlikely to get sick from COVID-19.
5	[16]	Healthcare workers	Questionnaire	NA	(75.6%) of respondents said that COVID-19 is a seriously dangerous and 69.3% perceived that they are at high risk of contracting the disease
6	[17]	Population	phone-based survey	NA	Majority believe that practicing social/physical Distancing makes difference in preventing contracting of the virus.
7	[18]	Students (Amhara)	Questionnaire	NA	230 (56.4%) [95% CI 51.2%, 61%] of college students had positive attitude in the prevention and control strategies of COVID-19 pandemic.
8	[19]	Nurses	Questionnaire	NA	The study participants had good attitude on COVID-19
9	[20]	Patients visiting JMC	Interview	NA	77.3% of visitors frequently washed their hands with water and soap. 90.3% avoided crowded place.
10	[23]	Southern Ethiopia population	Interview	Mean score 34.45 (±SD 5.5)	(90.3) had good attitude toward covid-19 and its prevention.

Table 4: Studies on practice.

No	Author	Participant	Instrument	Practice score	Result
1	[12]	Patients (chronic diseases)	Questioner	NA	47.3% (95% CI (42.4–52.2%)) had Poor practice. Only 105 (25.9%) of study participants had a good practice.
2	[13]	Arba Minch Town (Population)	Online questioner	NA	336(63.3%) were didn't used face-mask.
3	[14]	Students at Debre Berhan University	Questioner	NA	Overall high levels of attitude and good practice
4	[15]	Population	Telephone interview	NA	About 43% 265 of the respondents never practice any of the COVID-19 prevention methods and only less than 266 one fifth (19.1%) of the respondents follow COVID-19 prevention measures either usually or 267 always.
5	[16]	Healthcare workers	Questionnaire	NA	63.5% of the surveyed healthcare workers followed Correct practices regarding COVID-19.
6	[18]	Students (Amhara)	Questionnaire	NA	265 (65%) [95% CI 60, 70.1%] of college students had good level of Prevention practice.
7	[19]	Nurses	Questionnaire	NA	278(67%) had good prevention practice
8	[20]	Patients visiting JMC	Interview	NA	Good practice
9	[22]	Educated individuals	Online questioner	NA	(285/528, 54%) of the respondents had good COVID-19 preventive practice.
10	[21]	Population	Online questioner	3.09 ± 1.06 Range (0-6)	About 77.4% of the respondents were not obeying government restrictions
11	[23]	Southern Ethiopia population	Interview	NA	(80%) have bad practice and (93.3%) never used surgical mask.



CONCLUSION

In those thirteen studies on Knowledge, Attitude and Practice towards COVID-19 in Ethiopia; there is a gap between knowledge, attitude and practice; some had good knowledge about the virus but their practice was very poor that is why COVID-19 is increase alarmingly in Ethiopia. The review also revealed that knowledge directly influenced attitudes. Good KAP is a tool that can be used to assess the current conditions and if it is used properly it plays a pivotal role to control the spread of COVID-19. Therefore the following recommendations are given

- ✎ Government and ministry of health must give information to the society through TV, radio and social media repeatedly.
- ✎ Address peoples live in rural area through health professional about what is at stake
- ✎ Wearing mask must be mandatory
- ✎ Follow people's day to day activities when they are in the shop/malls, taxi, work etc.

DATA AVAILABILITY

All the datasets used to support the findings of this study are all in the text.

ACKNOWLEDGEMENTS

I am deeply grateful to all sources of materials used for this review article and it is duly acknowledged.

REFERENCE

1. Wang J, Wang Z. Strengths, weaknesses, opportunities and threats (SWOT) analysis of china's prevention and control strategy for the COVID-19 epidemic. *Int J Environ Res Public Health*. 2020; 17: 2235. DOI: 10.3390/ijerph17072235
2. Log S. Understanding SARS-CoV-2. 2020: 1-9.
3. Zhang C, Zheng W, Huang X, Bell EW, Zhou X, Zhang Y. Protein structure and sequence re-analysis of 2019-nCoV genome does not indicate snakes as its intermediate host or the unique similarity between its spike protein insertions and HIV-1. *J Proteome Res*. 2020. DOI: 10.1021/acs.jproteome.0c00129
4. Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID-19 infection: Origin, transmission, and characteristics of human coronaviruses. *J Adv Res*. 2020; 24: 91-98. DOI: 10.1016/j.jare.2020.03.005
5. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: A retrospective review of medical records. *Lancet*. 2020; 395: 809-815. DOI: 10.1016/S0140-6736(20)30360-3
6. Search, G. Coronavirus COVID-19 situation reports-World Health Organization. 2020; 5-7.
7. Camilla Rothe, Mirjam Schunk, Peter Sothmann, Gisela Bretzel, Guenter Froeschl, Claudia Wallrauch, et al. Transmission of 2019- nCoV infection from an asymptomatic contact in Germany. *N Engl J Med*. 2020; 382: 970-971. DOI: 10.1056/NEJMc2001468
8. Chaolin Huang, Yeming Wang, Xingwang Li, Lili Ren, Jianping Zhao, Yi Hu, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020; 395: 497-506. DOI: 10.1016/S0140-6736(20)30183-5
9. WHO. Coronavirus disease 2019 (COVID-19) Situation Report-195. WHO, 2020.
10. Ethiopian Public Health Institute. COVID-19 Interim containment strategies-Ethiopia. Addis Ababa, March, 2020.
11. Haq N, Hassali MA, Shafe AA, Saleem F, Farooqui M, Aljadhey H. A cross sectional assessment of knowledge, attitude and practice towards Hepatitis B among healthy population of Quetta, Pakistan. *BMC Public Health*. 2012; 12: 692. DOI: 10.1186/1471-2458-12-692
12. Yonas Akalu, Birhanu Ayelign, Meseret Derbew Molla. Knowledge, attitude and practice towards covid-19 among chronic disease patients at Addis Zemen Hospital, Northwest Ethiopia. *Infect Drug Resist*. 2020; 13: 1949-1960. DOI: 10.2147/IDR.S258736
13. Tewodros Fantaye Nigussie, Nuredin Nassir Azmach. Knowledge, attitude and practice towards covid-19 among Arba Minch town, southern Ethiopia. *Global Scientific Journal*: 2020; 8. <https://tinyurl.com/y69qsdzc>
14. Yared Asmare Aynalem, Tadesse Yirga Akalu, Birhan Gebresellassie, Nigussie Tadesse Sharew, Wondimeneh Shibabaw Shiferaw. Assessment of undergraduate student knowledge, practices, and attitude towards COVID-19 in Debre Berhan University, Ethiopia. 2020. DOI: 10.21203/rs.3.rs-28556/v1
15. Edessa Negera, Tesfaye Moti Demissie, Ketema Tafess. Inadequate level of knowledge, mixed outlook and poor adherence to COVID-19 prevention guideline among Ethiopians.2020. DOI: 10.1101/2020.07.22.215590
16. Bedru Jemal, Zemedu Aweke Ferede, Simeneh Mola, Sileshi Hailu et al. Knowledge, attitude and practice of healthcare workers towards COVID-19 and its prevention in Ethiopia: A multicenter study. 2020. DOI: 10.21203/rs.3.rs-29437/v1
17. Nigatu R Geda, Belaineh Legesse, Sindu W Kebede, Belay Fekadu. Knowledge and attitude towards covid-19 in Ethiopia. *Frontieri*. 2020.
18. Abay Woday Tadesse, Negesse Melese, Sisay Eshetie, Mulugeta Chanie, Ahmed Ali. Knowledge, attitude, and practice and associated factors towards COVID-19 among college students in Amhara Region, Ethiopia; A cross-sectional study. 2020. DOI: 10.21203/rs.3.rs-38099/v1
19. Degenah Bahrey, Gebremeskel Tukue, Gebre Teklemariam. Knowledge, Attitude, practice and psychological response toward COVID-19 among nurses during the COVID-19 outbreak in Northern Ethiopia, 2020. DOI: 10.21203/rs.3.rs-26236/v1.
20. Yohannes Kebede, Yimenu Yitayih, Zewdie Birhanu, Seblework Mekonen, Argaw Ambelu. Knowledge, perceptions and preventive practices towards COVID-19 early in the outbreak among Jimma University medical center visitors, Southwest Ethiopia. *PLoS One*. 2020; 15: e0233744. DOI: 10.1371/journal.pone.0233744
21. Daniel Bekele, Tadesse Tolossa, Reta Tsegaye, Wondesen Teshome. The knowledge and practice towards COVID-19 pandemic prevention among residents of Ethiopia: An online cross-sectional study. 2020. DOI: 10.1101/2020.06.01.127381
22. Henok Dagne, Kassahun Alemu, Baye Dagnaw, Demewoz Tadesse, Abiy Maru Alemayehu, Zewudu Andualem, et al. Prevention practice and associated factors of coronavirus disease 2019 (COVID-19) outbreak among educated Ethiopians: An online based cross-sectional survey. 2020. DOI: 10.21203/rs.3.rs-34504/v1
23. Simeneh Mola, Zemedu Aweke, Bedru Jemal, Robel Hussein, Sileshi Hailu, Derartu Neme, et al. Magnitude and associated factors for attitude and practice of Southern Ethiopian residents toward COVID-19 and its preventions: A community based cross sectional study. 2020. DOI: 10.21203/rs.3.rs-36120/v1
24. Biruk Wogayehu, Wondimagegn Taye, Yilma Chisha, Kahssay Faraja. Knowledge regarding 2019 novel coronavirus (2019-nCoV) infection among final year health science students at Arbaminch College of Health Sciences, Southern Ethiopia: A cross-sectional study. 2020. DOI: 10.21203/rs.3.rs-24777/v1